

Notre-Dame Lifesciences



PRODUCTS

BRINGING INNOVATION TO LIFE

Toll Free
1800-532-6700

www.notredame.co.in

info@notredame.co.in

Pantodame[™] **40**

Pantoprazole 40 mg

Pantodame[™] **DSR 40**

Pantoprazole (40 mg) + Domperidone (30 mg)

EsmoDame[™] **Fast 40**

Esomeprazole 40 mg

Esmodame[™] **DSR 40**

Esomeprazole (40 mg) + Domperidone (30 mg)

DameFlex[™] SP

**Aceclofenac (100 mg) + Paracetamol (325 mg)
+ Serratiopeptidase (15 mg)**

DameFlex[™] DCP


**Diclofenac Sodium 30 mg + Chiorzorazone 500 mg Tablets
+ Paracetamol 325 mg**

Damekast[™] Liv

Montelukast (10 mg) + Levocetirizine (5 mg)

**Reliable relief,
Right on schedule.**

Pantodame[™] 40



Pantoprazole 40 mg

Indications

- Acidity and acid-related dyspepsia
- Treatment of heartburn/GERD
- Peptic ulcer disease
- Zollinger–Ellison syndrome
- Prophylaxis of NSAID-induced ulcers (as directed by physician)

Storage

- Store below 30°C.
- Keep in the original blister, protected from moisture and light.
- Keep out of reach of children.

Pantodame-40 is a proton pump inhibitor (PPI) that effectively reduces excess stomach acid. It is indicated for:

- Heartburn and gastroesophageal reflux disease (GERD/acid reflux)
- Peptic ulcer disease (gastric and duodenal ulcers)
- Zollinger–Ellison syndrome (pathological hypersecretion)
- Prevention of NSAID-associated gastric/duodenal ulcers (as prescribed)

By lowering gastric acid secretion, Pantodame-40 relieves symptoms such as burning pain, regurgitation, and acidity while supporting mucosal healing.

How It Works

Pantoprazole selectively and irreversibly inhibits the H⁺/K⁺ ATPase proton pump in gastric parietal cells, the final step in acid production. This provides potent and sustained suppression of basal and stimulated gastric acid secretion.

Key Benefits

- Potent acid suppression for rapid relief from heartburn and epigastric pain
- Promotes healing of erosions and ulcers
- Reduces recurrence of reflux symptoms with continued therapy as prescribed
- Once-daily convenience for most indications

Directions for Use

- Take exactly as prescribed by your physician.
- Administer on an empty stomach, ideally 30–60 minutes before breakfast.
- If a second daily dose is prescribed, take it before the evening meal.
- Swallow the tablet whole with water; do not split, crush, or chew.
- Do not exceed the recommended dose or duration. Complete the full course even if symptoms improve early.

Safety Advice

Alcohol: Avoid; alcohol can worsen gastric irritation and reflux.

Pregnancy: Consult your doctor. Use only if potential benefits outweigh risks.

Breastfeeding: Likely compatible if prescribed; discuss with your physician.

Driving/Operating Machines: Caution if you experience dizziness or visual disturbance.

Kidney: Generally safe; no routine dose adjustment required.

Liver: Use with caution in severe hepatic impairment; dose adjustment may be required.

Common Side Effects

Usually mild and transient. Consult your doctor if persistent or troublesome.

- Headache, dizziness
- Nausea, vomiting
- Abdominal pain, flatulence, diarrhea
- Upper respiratory tract infection
- Joint pain

Long-term/high-dose use (especially >1 year) may increase the risk of:

- Bone fractures (hip, wrist, spine)
- Vitamin B12 deficiency
- Low magnesium (hypomagnesemia)

Seek urgent care for severe abdominal pain, black/tarry stools, blood in vomit, persistent diarrhea, or allergic reactions.

Don't let
ACID PEPTIC DISORDERS
win over the stomach

**Sure Shot
Win over APD**



Pantodame[™] DSR 40

Pantoprazole (40 mg) + Domperidone (30 mg)

Indications

- **Dyspepsia**
- **Hyperacidity**
- **Gastritis**
- **Nausea & Vomiting**
- **Heartburn**

Storage

- Store below 30°C.
- Keep in the original pack, protected from moisture and light.
- Keep out of reach of children.

Long Lasting Relief from Acid Peptic Disorders

Pantodame-DSR 40 is a dual-action gastroprokinetic and acid-suppressing capsule used for the management of:

- **Gastroesophageal Reflux Disease (GERD/acid reflux)**
- **Dyspepsia (indigestion)**
- **Gastritis**

It reduces excess stomach acid, relieves heartburn and epigastric pain, and improves gastric motility to ease bloating and discomfort.

How It Works

- Pantoprazole: Proton pump inhibitor that blocks gastric acid secretion, relieving heartburn and acid-related irritation.
- Domperidone: Prokinetic that increases movement of the stomach and intestines, helping food pass more easily and reducing nausea, bloating, and early satiety.

Key Benefits

- Rapid relief from heartburn and acid-related pain
- Reduces gastric acidity and reflux episodes
- Improves gastric emptying and decreases bloating
- Supports symptom control for better daily comfort

Directions for Use

- Take exactly as prescribed by your physician.
- Swallow whole; do not chew, crush, or open the capsule.
- Take on an empty stomach, ideally 30–60 minutes before breakfast. If a second daily dose is prescribed, take it before the evening meal.
- Use the lowest effective dose for the shortest duration as advised.

Common Side Effects

Usually mild and transient. Consult your doctor if persistent or troublesome.

- Diarrhea
- Stomach pain
- Flatulence
- Dry mouth
- Dizziness or drowsiness
- Headache
- Nausea/vomiting

Do not drive or operate machinery if you feel dizzy or sleepy.

Safety Advice


Alcohol: Use with caution. Alcohol may worsen gastric irritation and drowsiness.
Pregnancy: Consult your doctor. Use only if potential benefits outweigh risks.
Breastfeeding: Use with caution; your doctor may advise timing feeds or temporary interruption.
Driving/Operating Machines: Avoid if drowsy, dizzy, or with blurred vision.
Kidney: Use with caution; dose adjustment may be needed—consult your doctor.
Liver: Generally safe if prescribed; your doctor will advise based on your liver function.

Pantodame[™] DSR 40

Turn down ↓ the acid,
Turn up ↑ the comfort.

EsmoDame[®] Fast 40

Esomeprazole 40 mg



Indications

- Gastroesophageal Reflux Disease (GERD), including erosive esophagitis
- Symptomatic acid reflux/heartburn
- Peptic ulcer disease (gastric and duodenal)
- Zollinger–Ellison syndrome (pathological hypersecretion)
- Prevention of NSAID-associated gastric/duodenal ulcers (as prescribed)

Storage

- Store below 30°C.
- Keep in the original blister to protect from moisture and light.
- Keep out of reach of children.

EsmoDame-Fast 40 is a proton pump inhibitor (PPI) that suppresses excessive gastric acid secretion. By reducing stomach acid, it relieves heartburn, epigastric pain, and acid regurgitation and supports mucosal healing.

How It Works

Esomeprazole selectively and irreversibly inhibits the H⁺/K⁺ ATPase (“proton pump”) in gastric parietal cells—the final step of acid production—resulting in potent and sustained suppression of basal and stimulated gastric acid secretion.

Key Benefits

- Rapid relief from heartburn and acid-related discomfort
- Promotes healing of erosive esophagitis and peptic ulcers
- Helps reduce recurrence of reflux symptoms with continued therapy (as advised)
- Once-daily convenience for most indications

Directions for Use

- Take exactly as prescribed by your physician.
- Administer on an empty stomach, ideally 30–60 minutes before breakfast.
- If a second daily dose is prescribed, take it before the evening meal.
- Swallow the tablet whole with water; do not split, crush, or chew.
- Complete the full course even if symptoms improve early. Do not exceed the recommended dose or duration.

Common Side Effects

Usually mild and transient. Consult your doctor if persistent or troublesome.

- Headache
- Nausea, vomiting
- Diarrhea or constipation
- Abdominal pain, flatulence
- Long-term/high-dose use (especially >1 year) may increase the risk of:
Bone fractures (hip, wrist, spine)
Vitamin B12 deficiency
Low magnesium (hypomagnesemia)

Seek urgent care for black/tarry stools, blood in vomit, severe abdominal pain, persistent diarrhea, or signs of allergy.

Safety Advice

Alcohol: Use with caution; alcohol can aggravate reflux and gastric irritation.

Pregnancy: Consult your doctor. Use only if potential benefits outweigh risks.

Breastfeeding: Use with caution; your doctor may advise timing feeds or alternative options.

Driving/Operating Machines: Avoid if you experience dizziness, sleepiness, or blurred vision.

Kidney: Generally safe; no routine dose adjustment required. Use caution in severe renal insufficiency.

Liver: Use with caution in severe hepatic impairment; your doctor may adjust the dose.



EsmodameTM DSR 40

Esomeprazole (40 mg) + Domperidone (30 mg)

- More effective than Omeprazole in healing and
- symptom resolution in GERD patients with reflux esophagitis
- Provides greater acid control & maintains
- intra-gastric pH greater than 4 for a longer period

Indications

- Treatment of GERD (acid reflux)
- Symptomatic relief of acid-induced epigastric pain / heartburn
- Management of acid-related dyspepsia
- Adjunct in gastritis with reflux symptoms (as prescribed)

Esmodame-DSR 40 is a sustained-release dual-action formulation for the management of acid-related upper gastrointestinal disorders. It combines a prokinetic (Domperidone) to enhance gastric motility with a proton pump inhibitor (Esomeprazole) to suppress excess acid.

This synergy helps relieve:

- Gastroesophageal Reflux Disease (GERD / Acid Reflux)
- Persistent heartburn and epigastric pain
- Acid-related dyspeptic symptoms (bloating, early fullness)
- Acid-associated gastritis discomfort

How It Works

- **Domperidone:** Enhances coordinated gastric emptying by increasing motility in the upper GI tract; reduces nausea, bloating, and early satiety.
- **Esomeprazole:** Irreversibly inhibits the H⁺/K⁺ ATPase (proton pump) in gastric parietal cells, producing potent and sustained reduction of gastric acid secretion.

Key Benefits

- **Dual relief:** reduces acid AND improves gastric motility
- Rapid symptom control for burning, regurgitation, and sour taste
- Decreases post-meal fullness and upper abdominal discomfort
- Supports mucosal healing when used as prescribed
- Enhances quality of life in chronic reflux patients

Directions for Use

- Take exactly as prescribed by your physician.
- Administer on an empty stomach: ideally 30–60 minutes before the first meal of the day.
- Swallow capsule whole; do not open, chew, or crush.
- If a second daily dose is prescribed, take before the evening meal (only if directed).
- Do not exceed the prescribed duration or dose.
- Continue therapy for the full recommended course even if symptoms improve early.

Safety Advice

Alcohol: Use cautiously; may aggravate gastric irritation and increase dizziness.

Pregnancy: Consult your doctor. Use only if benefits outweigh potential risks.

Breastfeeding: Not recommended; Domperidone may pass into breast milk—avoid unless specifically advised.

Driving / Operating Machinery: Avoid if drowsy, dizzy, or experiencing visual disturbance.

Kidney: Use with caution in severe renal impairment; medical supervision required.

Liver: Generally safe if prescribed; inform your doctor of any existing liver disease.

Common Side Effects

(Usually Mild & Transient)

- Diarrhea
- Flatulence
- Stomach / abdominal pain
- Dry mouth
- Headache
- Dizziness or drowsiness
- Nausea

Seek medical advice if symptoms are persistent, severe, or unusual.



DameFlexTM SP

**Aceclofenac (100 mg) + Paracetamol (325 mg)
+ Serratiopeptidase (15 mg)**

Indications

- Pain relief in musculoskeletal and joint conditions
- Inflammatory arthropathies (e.g., RA, OA, ankylosing spondylitis)
- Post-operative and post-traumatic pain and edema
- Dental pain (as prescribed)

DameFlex- SP is a triple-action analgesic and anti-inflammatory tablet formulated for short-term relief of pain, swelling, and stiffness affecting muscles and joints. It is beneficial in conditions such as osteoarthritis, rheumatoid arthritis, ankylosing spondylitis, musculoskeletal injuries, dental pain, and post-operative pain.

- Targets pain and fever
- Reduces inflammation and tissue swelling
- Supports faster functional recovery

How It Works

- **Aceclofenac:** A non-steroidal anti-inflammatory drug (NSAID) that inhibits prostaglandin synthesis to reduce pain and inflammation.
- **Serratiopeptidase:** A proteolytic enzyme that helps break down inflammatory proteins at the site of injury, decreasing edema and promoting healing.
- **Paracetamol:** An analgesic and antipyretic that helps relieve pain and reduce fever via central mechanisms.

Key Benefits

- Comprehensive control of pain, swelling, and stiffness
- Helps restore mobility and daily activity comfort
- Faster resolution of post-injury and post-surgical inflammation

Common Side Effects

Usually mild and transient. Consult your doctor if persistent or troublesome.

- Nausea, vomiting, diarrhea
- Stomach/abdominal pain, indigestion, heartburn, loss of appetite
- Dizziness, drowsiness
- Elevated liver enzymes
- Headache

Seek immediate medical attention for black/tarry stools, vomiting blood, severe abdominal pain, yellowing of skin/eyes, or unusual bruising/bleeding.

Directions for Use

- Take exactly as prescribed by your physician.
- Swallow whole with water; do not chew, crush, or break.
- Take with food or milk to minimize stomach upset.
- Use the lowest effective dose for the shortest duration recommended.
- Do not exceed the prescribed dose or frequency.

Safety Advice

- **Alcohol:** Do not consume alcohol while taking DameFlex-SP; it increases the risk of stomach bleeding and liver toxicity.
- **Pregnancy:** Consult your doctor. Use may be unsafe during pregnancy; prescribed only if benefits outweigh risks.
- **Breastfeeding:** Safety data are limited—use only on medical advice.
- **Driving/Operating Machinery:** Avoid if you feel dizzy, sleepy, or your vision is affected.
- **Kidney:** Use with caution; dose adjustment may be required. Avoid in severe renal impairment.
- **Liver:** Use with caution; dose adjustment may be required. Do not use in severe or active liver disease.

For

SPEEDY RELIEF

in Painful Musculo-Skeletal Disorders

The Powerful Trio of

Anti-inflammatory Analgesic & Musculo-Skeletal Relaxant

DameFlex™ DCP

Diclofenac Sodium 30 mg + Chiorzorazone 500 mg Tablets + Paracetamol 325 mg



Diclofenac

- Offers strong & selective anti-inflammatory action
- Better in acute inflammatory conditions for quick relief

Chiorzoxazone

- Relaxes skeletal muscles & relieves spasm
- Offers central, safe & selective pain relief

Paracetamol

- Ensures faster analgesia
- Synergistic pain relieving action

Storage

- Store below 30°C.
- Keep in the original package, protected from moisture and light.
- Keep out of reach of children.

- Lower Back Pain
- Spondylitis
- Frozen Shoulder
- Tendonitis
- Dislocations
- Sports Injuries

Dameflex-DCP Tablet is a triple-action formulation used for the relief of pain associated with acute muscle spasms. It helps reduce muscle stiffness, improves mobility, and alleviates discomfort so you can return to daily activities more comfortably

- Centrally acting muscle relaxant + dualanalgesic/anti-inflammatory combination
- Targets pain, inflammation, and spasm to improve range of motion

Uses

Treatment of pain due to muscle spasm

How Dameflex-DCP Works ?

- Chlorzoxazone: Acts on the brain and spinal cord centers to relieve muscle stiffness and spasms.
- Diclofenac and Paracetamol: Reduce pain and inflammation by blocking chemical messengers responsible for pain, redness, and swelling.

Benefits

- Relieves painful muscle spasms
- Reduces inflammation and swelling
- Eases stiffness to improve muscle movement
- Supports faster return to normal activities

How to Use

- Take exactly as prescribed by your doctor.
- Swallow whole; do not chew, crush, or break.
- May be taken with or without food; taking with food can help reduce stomach upset.
- Use the lowest effective dose for the shortest duration as advised.

DameFlex™ DCP

Common Side Effects

Most side effects are mild and resolve as your body adjusts. Contact your doctor if they persist or bother you.

- Dizziness
- Nausea
- Indigestion
- Abdominal pain
- Constipation

Kidney: Not recommended in kidney disease; consult your doctor.
Liver: Use with caution in liver disease; dose adjustment may be required. Consult your doctor.

Safety Advice

Alcohol: Consult your doctor. Safety with alcohol is not established.

Pregnancy: Consult your doctor. May be unsafe during pregnancy; your doctor will weigh risks and benefits.

Breastfeeding: Consult your doctor. May pass into breast milk and could harm the baby.

Driving/Operating Machines: Unsafe if drowsy or dizzy. Dameflex-DCP may decrease alertness or affect vision.

The Novel & Effective Approach in Treatment of Allergy & Asthma

Selectively Treats Seasonal & Persistent Allergic Rhinitis

Damekast[™] Liv

Montelukast (10 mg) + Levocetirizine (5 mg)

Damekast-Liv is a dual-action anti-allergic tablet formulated to relieve symptoms of allergic rhinitis and related conditions, including:

- Sneezing, runny/stuffy nose
- Itching, swelling
- Watery eyes, congestion

It reduces airway inflammation and helps make breathing easier.



Indications

- Treatment of sneezing and runny nose due to allergies
- Treatment of hay fever (allergic rhinitis)
- Treatment of allergic skin conditions with redness and itching (as prescribed)

How It Works

- **Levocetirizine:** A non-sedating antihistamine that blocks histamine (the chemical responsible for sneezing, watery eyes, and itching).
- **Montelukast:** A leukotriene receptor antagonist that blocks leukotrienes, reducing inflammation and swelling in the airways and nose.

Key Benefits

- Comprehensive relief from nasal and ocular allergy symptoms
- Reduces itching, redness, and skin irritation in allergic skin conditions
- Helps ease airway inflammation to support easier breathing
- Once-daily convenience (as directed by your physician)

Common Side Effects

Usually mild and temporary. Consult your doctor if persistent or troublesome.

- Nausea, vomiting, diarrhea
- Dry mouth, headache, fatigue, sleepiness, dizziness
- Skin rash
- Upper respiratory tract infection, sore throat, fever
- Abdominal pain
- Increased liver enzymes (rare; monitor if advised)

Safety Advice

- **Alcohol:** Use with caution; may increase drowsiness. Consult your doctor.
- **Pregnancy:** Generally considered safe if prescribed. Follow medical advice.
- **Breastfeeding:** Not recommended; consult your doctor before use.
- **Driving/Operating Machines:** Avoid if drowsy, dizzy, or vision is affected.
- **Kidney:** Use with caution in kidney disease. Not recommended in severe renal impairment; your doctor may adjust the dose.
- **Liver:** Use with caution in severe liver disease. Limited data in severe impairment; no adjustment typically needed in mild-to-moderate disease—follow your doctor’s advice.

Avoid driving or operating machinery if you feel sleepy, dizzy, or have blurred vision.



Polar Frost ICE SPRAY is a ready-to-use ice spray can. Providing soothing pain relief, the ice spray is equipped with a special dispenser valve that directs the flow to the area to be treated.

- **Fast, Targeted Pain Relief**
- **Easy Use Allows Swift And Direct Pain Relief**
- **Fast Acting Pain Relief For Strains, Sprains, Muscles And Joints**
- **Rapidly Cools The Injured Area**
- **Ideal For Areas That Are Hard To Reach Or Rub**

Used for relieving pain caused by injuries, scrapes, falls or sprains, and in cases requiring cryotherapy and application of cold. Thanks to its anaesthetic effect induced by cold on the injured area, sports activities can be resumed quickly.

TECHNICAL DATA SHEET

For optimal use, consult a healthcare professional before applying to children and the elderly. Suitable for individuals over the age of 3.

Ingredients: Gas: Isobutane, Buthane, Propane, Menthol
Shelf Life : 5 Years From Production Date.
Storage: Keep At Ambient Temperature, In A Dry Place, Away From Heat Sources, Flames Or Sparks
Packaging: Box Of 24 Cans
Classification: Medical Device Class Iia Classification According To The Annex Ix Paragraph Iii Rule 9 Of The European Directive 93/42/Eec
Dimension: MI 200, MI 400
Usage : For optimal use, consult a healthcare professional before applying to children and the elderly. Suitable for individuals over the age of 3. For external use only.
Manufactured by : Dispotech Srl - Via al piano 29 23020 Gordona (SO), Italy
Marketed by : Notre-DameLifesciences Private Limited



Keep it at a distance of at least 20 cm from the affected part during application to avoid skin irritation or mild burns.

Listening. Innovating.

Transforming Healthcare: Your Needs, Our Inventions

At Notre-Dame Lifesciences, we believe that groundbreaking innovation stems from a deep understanding of our customers' needs. Our approach is rooted in active listening and engagement with healthcare professionals, patients, and partners across the globe. By carefully analyzing feedback, market trends, and emerging healthcare challenges, we identify gaps in current offerings and opportunities for improvement. This customer-centric philosophy drives our research and development team to transform insights into cutting-edge solutions that address real-world healthcare needs.

About us

Mission

To revolutionize healthcare by developing innovative, high-quality, and affordable pharmaceutical, medical device, and cosmetic solutions. We are committed to listening intently to our customers, transforming their needs into groundbreaking inventions, and delivering products that improve health outcomes and enhance quality of life for people around the world.

Vision

To be a global leader in customer-driven healthcare innovation, recognized for our ability to anticipate and address emerging medical challenges. We envision a future where our inventions set new industry standards, making advanced healthcare solutions accessible to all, and positioning Notre-Dame Lifesciences as the preferred partner for healthcare professionals and patients worldwide.

Notre-Dame Lifesciences Private Limited

50/60, A4, Kulangara Arcade, Varissery Junction, Mariathuruthu P
Kottayam, Kerala, Pin 686017

Phone number : +91 8891226474, +91 8943169662



Toll Free
1800-532-6700

Thank you

www.notredame.co.in

www.polarfrostice.com